

Global Flow


Anandan

Affine

Global Motion

- Estimate motion using all pixels in the image.
- Parametric flow gives an equation, which describes optical flow for each pixel.
 - Affine
 - Projective
- Global motion can be used to
 - generate mosaics
 - Object-based segmentation

Affine



$$u(x, y) = a_1x + a_2y + b_1$$

$$v(x, y) = a_3x + a_4y + b_2$$

$$U = X - X'$$

$$\begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} x' \\ y' \end{bmatrix}$$


Affine

$$u(x, y) = a_1x + a_2y + b_1$$

$$v(x, y) = a_3x + a_4y + b_2$$

$$\begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

Spatial Transformations




translation



rotation



shear



Rigid (rotation and translation)



affine

Anandan

$$u(x, y) = a_1x + a_2y + b_1$$

$$v(x, y) = a_3x + a_4y + b_2$$

•Affine

$$\begin{bmatrix} u(x, y) \\ v(x, y) \end{bmatrix} = \begin{bmatrix} x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ b_1 \\ a_3 \\ a_4 \\ b_2 \end{bmatrix}$$

$$\mathbf{u}(\mathbf{x}) = \mathbf{X}(\mathbf{x})\mathbf{a}$$

Anandan

$$\mathbf{u}(\mathbf{x}) = \mathbf{X}(\mathbf{x})\mathbf{a}$$

Optical flow constraint eq

$$f_x u + f_y v = -f_t$$

$$E(\mathbf{u}) = \sum_{\forall \mathbf{x} \in f(x,y)} (f_t + f_x^T \mathbf{u})^2$$

$$E(\mathbf{a}) = \sum_{\forall \mathbf{x} \in f(x,y)} (f_t + f_x^T \mathbf{X}(\mathbf{x})\mathbf{a})^2$$

$$f_x = \begin{bmatrix} f_x \\ f_y \end{bmatrix}$$

$$E(\delta \mathbf{a}) = \sum_{\forall \mathbf{x} \in f(x,y)} (f_t + f_x^T \mathbf{X} \delta \mathbf{a})^2$$

(a) Derive this Due

Anandan

$$E(\delta a) = \sum_{\forall x \in f(x,y)} (f_t + f_x^T X \delta a)^2$$

min 

$$\left[\sum X^T (f_x) (f_x)^T X \right] \delta a = - \sum X^T f_x f_t \quad \text{(a) Derive this}$$

$$Ax = b$$

Linear system

Basic Components

- Pyramid construction
- Motion estimation
- Image warping
- Coarse-to-fine refinement




Image Warping

- Warping an image f into image h using some transformation g , involves mapping intensity at each pixel (x,y) in image f to a pixel $(g(x),g(y))$ in image h such that

$$(x', y') = (g(x), g(y))$$

- In case of affine transformation, $x = (x, y)$ is transformed to $x' = (x', y')$ as:

$$\mathbf{x}' = \mathbf{Ax} + \mathbf{b}$$

Image Warping




Image Warping

$$X' = X - U = X - (AX + b)$$

$$X' = (I - A)X - b$$

$$X' = A'X - b$$

$$X' + b = A'X$$

$$(A')^{-1}(X' + b) = X$$



$$(A')^{-1}(X' + b) = X'' \quad X' \rightarrow X''$$

Image at time t: X
Image at time t-1: X'

Image Warping

- How about values in $X'' = (x'', y'')$ are not integer.
- But image is sampled only at integer rows and columns
 - Instead of converting X' to X'' and copying $f(X', t-1)$ at $f(X'', t-1)$ we can convert integer values X'' to X' and copy $f(X', t-1)$ at $f(X'', t-1)$

Image Warping




Image Warping

- But how about the values in X' are not integer.
- Perform bilinear interpolation to compute $f(X', t-1)$ at non-integer values.


Image Warping

$$(A')^{-1}(X' + b) = X''$$

$$(X' + b) = (A')X''$$

$$X' = (A')X'' - b \quad X'' \rightarrow X'$$

Warping



Video Mosaic




Video Mosaic



Video Mosaic




Sprite




Szeliski

Projective

Projective



$f(X', t-1)$



$f(X, t)$

$$x' = \frac{a_1x + a_2y + b_1}{c_1x + c_2y + 1}$$

$$y' = \frac{a_3x + a_4y + b_2}{c_1x + c_2y + 1}$$

Szeliski

$$x' = \frac{a_1x + a_2y + b_1}{c_1x + c_2y + 1} \quad \text{Projective}$$

$$y' = \frac{a_3x + a_4y + b_2}{c_1x + c_2y + 1}$$

$$E = \sum [f(x', y') - f(x, y)]^2 = \sum e^2$$

↓ min

Szeliski

Motion Vector:

$$\mathbf{m} = [a_1 \quad a_2 \quad a_3 \quad a_4 \quad b_1 \quad b_2 \quad c_1 \quad c_2]^T$$

Szeliski (Levenberg-Marquadet)

$$\alpha_{kl} = \sum_n \frac{\partial e_n}{\partial m_k} \frac{\partial e_n}{\partial m_l}$$

$$b_k = -\sum e \frac{\partial e_n}{\partial m_k}$$

gradient

$$\Delta \mathbf{m} = (\mathbf{A} + \lambda \mathbf{I})^{-1} \mathbf{b}$$

Approximation of
Hessian ($J^T J$, Jacobian)

$$\frac{\partial e}{\partial a_1} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_1} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_1}$$

$$E = \sum [f(x', y') - f(x, y)]^2 = \sum e^2$$

$$x' = \frac{a_1 x + a_2 y + b_1}{c_1 x + c_2 y + 1}$$

$$y' = \frac{a_3 x + a_4 y + b_2}{c_1 x + c_2 y + 1}$$

Approximation of Hessian

$$J^T = \begin{bmatrix} \frac{\partial e_1}{\partial m_1} & \frac{\partial e_2}{\partial m_1} & \dots & \frac{\partial e_n}{\partial m_1} \\ \frac{\partial e_1}{\partial m_2} & \frac{\partial e_2}{\partial m_2} & \dots & \frac{\partial e_n}{\partial m_2} \\ \frac{\partial e_1}{\partial m_3} & \frac{\partial e_2}{\partial m_3} & \dots & \frac{\partial e_n}{\partial m_3} \\ \frac{\partial e_1}{\partial m_4} & \frac{\partial e_2}{\partial m_4} & \dots & \frac{\partial e_n}{\partial m_4} \\ \frac{\partial e_1}{\partial m_5} & \frac{\partial e_2}{\partial m_5} & \dots & \frac{\partial e_n}{\partial m_5} \\ \frac{\partial e_1}{\partial m_6} & \frac{\partial e_2}{\partial m_6} & \dots & \frac{\partial e_n}{\partial m_6} \\ \frac{\partial e_1}{\partial m_7} & \frac{\partial e_2}{\partial m_7} & \dots & \frac{\partial e_n}{\partial m_7} \\ \frac{\partial e_1}{\partial m_8} & \frac{\partial e_2}{\partial m_8} & \dots & \frac{\partial e_n}{\partial m_8} \end{bmatrix} \quad J = \begin{bmatrix} \frac{\partial e_1}{\partial m_1} & \frac{\partial e_1}{\partial m_2} & \frac{\partial e_1}{\partial m_3} & \frac{\partial e_1}{\partial m_4} & \frac{\partial e_1}{\partial m_5} & \frac{\partial e_1}{\partial m_6} & \frac{\partial e_1}{\partial m_7} & \frac{\partial e_1}{\partial m_8} \\ \frac{\partial e_2}{\partial m_1} & \frac{\partial e_2}{\partial m_2} & \frac{\partial e_2}{\partial m_3} & \frac{\partial e_2}{\partial m_4} & \frac{\partial e_2}{\partial m_5} & \frac{\partial e_2}{\partial m_6} & \frac{\partial e_2}{\partial m_7} & \frac{\partial e_2}{\partial m_8} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{\partial e_n}{\partial m_1} & \frac{\partial e_n}{\partial m_2} & \frac{\partial e_n}{\partial m_3} & \frac{\partial e_n}{\partial m_4} & \frac{\partial e_n}{\partial m_5} & \frac{\partial e_n}{\partial m_6} & \frac{\partial e_n}{\partial m_7} & \frac{\partial e_n}{\partial m_8} \end{bmatrix}$$

$$A = J^T J$$

$$\alpha_{kl} = \sum_n \frac{\partial e_n}{\partial m_k} \frac{\partial e_n}{\partial m_l} \quad \text{A Matrix}$$

Gradient Vector

$$b = \begin{bmatrix} -\sum_n e_n \frac{\partial e_n}{\partial a_1} \\ -\sum_n e_n \frac{\partial e_n}{\partial a_2} \\ -\sum_n e_n \frac{\partial e_n}{\partial a_3} \\ -\sum_n e_n \frac{\partial e_n}{\partial a_4} \\ -\sum_n e_n \frac{\partial e_n}{\partial b_1} \\ -\sum_n e_n \frac{\partial e_n}{\partial b_2} \\ -\sum_n e_n \frac{\partial e_n}{\partial c_1} \\ -\sum_n e_n \frac{\partial e_n}{\partial c_2} \end{bmatrix}$$

Partial Derivatives wrt motion parameters

$$\frac{\partial e}{\partial a_1} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_1} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_1}$$

$$x' = \frac{a_1 x + a_2 y + b_1}{c_1 x + c_2 y + 1}, \quad y' = \frac{a_3 x + a_4 y + b_2}{c_1 x + c_2 y + 1}$$

$\frac{\partial x'}{\partial a_1} = \frac{x}{c_1 x + c_2 y + 1}$	$\frac{\partial y'}{\partial a_1} = 0$
$\frac{\partial x'}{\partial a_2} = \frac{y}{c_1 x + c_2 y + 1}$	$\frac{\partial y'}{\partial a_2} = 0$
$\frac{\partial x'}{\partial a_3} = 0$	$\frac{\partial y'}{\partial a_3} = \frac{x}{c_1 x + c_2 y + 1}$
$\frac{\partial x'}{\partial a_4} = 0$	$\frac{\partial y'}{\partial a_4} = \frac{y}{c_1 x + c_2 y + 1}$
$\frac{\partial x'}{\partial b_1} = \frac{1}{c_1 x + c_2 y + 1}$	$\frac{\partial y'}{\partial b_1} = 0$
$\frac{\partial x'}{\partial b_2} = 0$	$\frac{\partial y'}{\partial b_2} = \frac{1}{c_1 x + c_2 y + 1}$
$\frac{\partial x'}{\partial c_1} = \frac{-x(a_1 x + a_2 y + b_1)}{(c_1 x + c_2 y + 1)^2}$	$\frac{\partial y'}{\partial c_1} = \frac{-x(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2}$
$\frac{\partial x'}{\partial c_2} = \frac{-y(a_1 x + a_2 y + b_1)}{(c_1 x + c_2 y + 1)^2}$	$\frac{\partial y'}{\partial c_2} = \frac{-y(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2}$

Partial derivatives wrt image coordinates

$$E = \sum [f(x', y') - f(x, y)]^2 = \sum e^2$$

$$\frac{\partial e}{\partial x'} = f'_x$$

$$\frac{\partial e}{\partial y'} = f'_y$$

Partial derivatives

$$\frac{\partial e}{\partial a_1} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_1} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_1} = f'_x \frac{x}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial a_2} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_2} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_2} = f'_x \frac{y}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial a_3} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_3} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_3} = f'_y \frac{x}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial a_4} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial a_4} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial a_4} = f'_y \frac{y}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial b_1} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial b_1} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial b_1} = f'_x \frac{1}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial b_2} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial b_2} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial b_2} = f'_y \frac{1}{c_1 x + c_2 y + 1}$$

$$\frac{\partial e}{\partial c_1} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial c_1} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial c_1} = f'_x \frac{-x(a_1 x + a_2 y + b_1)}{(c_1 x + c_2 y + 1)^2} + f'_y \frac{-x(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2}$$

$$\frac{\partial e}{\partial c_2} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial c_2} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial c_2} = f'_x \frac{-y(a_1 x + a_2 y + b_1)}{(c_1 x + c_2 y + 1)^2} + f'_y \frac{-y(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2}$$

Gradient Vector

$$\mathbf{b} = \begin{bmatrix} -\sum e f_{x'} \frac{x}{c_1 x + c_2 y + 1} \\ -\sum e f_{x'} \frac{y}{c_1 x + c_2 y + 1} \\ -\sum e f_{y'} \frac{x}{c_1 x + c_2 y + 1} \\ -\sum e f_{y'} \frac{y}{c_1 x + c_2 y + 1} \\ -\sum e f_{x'} \frac{1}{c_1 x + c_2 y + 1} \\ -\sum e f_{y'} \frac{1}{c_1 x + c_2 y + 1} \\ \sum e x \left[\frac{f_{x'}(a_1 x + a_2 y + b_1) + f_{y'}(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2} \right] \\ \sum e y \left[\frac{f_{x'}(a_1 x + a_2 y + b_1) + f_{y'}(a_3 x + a_4 y + b_2)}{(c_1 x + c_2 y + 1)^2} \right] \end{bmatrix}$$

Szeliski (Levenberg-Marquadt)

- Start with some initial value of m , and $\lambda = .001$
- For each pixel I at (x_i, y_i)
 - Compute (x', y') using projective transform.
 - Compute $e = f(x', y') - f(x, y)$
 - Compute $\frac{\partial e}{\partial m_k} = \frac{\partial e}{\partial x'} \frac{\partial x'}{\partial m_k} + \frac{\partial e}{\partial y'} \frac{\partial y'}{\partial m_k}$

Szeliski (Levenberg-Marquadet)

-Compute A and b

-Solve system

$$(A - \lambda I)\Delta m = b$$

-Update

$$m^{t+1} = m^t + \Delta m$$

Szeliski (Levenberg-Marquadet)

- check if error has decreased, if not increase λ by a factor of 10 and compute a new Δm
- If error has decreased, decrease λ by a factor of 10 and compute a new Δm
- Continue iteration until error is below threshold.