

Assignment # 8.1 Sample Key

1. Use reduction from **Halt** to show that one cannot decide **REPEATS**, where **REPEATS** = { f | for some x and y , $x \neq y$, $\varphi_f(x) \downarrow$, $\varphi_f(y) \downarrow$ and $\varphi_f(x) == \varphi_f(y)$ }

Let f, x be an arbitrary pair of natural numbers. $\langle f, x \rangle$ is in Halt iff $\varphi_f(x) \downarrow$

Define g by $\varphi_g(y) = \varphi_f(x) - \varphi_f(x)$, for all y .

Clearly, $\varphi_g(y) = 0$, for all y , iff $\varphi_f(x) \downarrow$, and $\varphi_g(y) \uparrow$, for all y , otherwise.

Summarizing, $\langle f, x \rangle$ is in Halt implies g is in REPEATS and $\langle f, x \rangle$ is not in Halt implies g is not in REPEATS

Halt \leq_m **REPEATS** as we were to show.

Note: I have not overloaded the index of a function with the function in my proof, but I do not mind if you do such overloading.

Assignment # 8.2 Sample Key

2. Show that **REPEATS** reduces to **Halt**. (1 plus 2 show they are equally hard)

Let f be an arbitrary natural number. f is in REPEATS iff for some x and y , $x \neq y$, $\varphi_f(x) \downarrow$, $\varphi_f(y) \downarrow$ and $\varphi_f(x) == \varphi_f(y)$

Define g by $\varphi_g(z) = \exists \langle x, y, t \rangle [STP(f, x, t) \ \& \ STP(f, y, t) \ \& \ (x \neq y) \ \& \ (VALUE(f, x, t) = VALUE(f, y, t))]$, for all z .

Clearly, $\varphi_g(z) = 1$, for all z , iff there is some pair, x, y , such that $\varphi_f(x) \downarrow$ and $\varphi_f(y) \downarrow$ and $\varphi_f(x) = \varphi_f(y)$, and $\varphi_g(z) \uparrow$, for all z , otherwise.

Summarizing, f is in REPEATS iff $\langle g, 0 \rangle$ is in Halt and so

REPEATS \leq_m **Halt** as we were to show.

Assignment # 8.3 Sample Key

3. Use Reduction from **Total** to show that **DOUBLES** is not even re, where

$$\text{DOUBLES} = \{ f \mid \text{for all } x, \varphi_f(x) \downarrow, \varphi_f(x+1) \downarrow \text{ and } \varphi_f(x+1) = 2 * \varphi_f(x) \}$$

Let f be an arbitrary natural number. f is in Total iff $\forall x \varphi_f(x) \downarrow$

Define g by $\varphi_g(x) = \varphi_f(x) - \varphi_f(x)$, for all x .

Clearly, $\varphi_g(x) = 0$, and so $\varphi_g(x+1) = 2 * \varphi_g(x) = 0$ for all x , iff $\forall x \varphi_f(x) \downarrow$; otherwise $\varphi_g(x) \uparrow$ for some x .

Summarizing, f is in Total iff g is in DOUBLES and so

TOTAL \leq_m **DOUBLES** as we were to show.

Assignment # 8.3 Alternate Key

3. Use Reduction from **Total** to show that **DOUBLES** is not even re, where

$$\text{DOUBLES} = \{ f \mid \text{for all } x, \varphi_f(x) \downarrow, \varphi_f(x+1) \downarrow \text{ and } \varphi_f(x+1) = 2 * \varphi_f(x) \}$$

Let f be an arbitrary natural number. f is in Total iff $\forall x \varphi_f(x) \downarrow$

Define g by $\varphi_g(x) = \varphi_f(x) - \varphi_f(x) + 2^x$ for all x .

Clearly, $\varphi_g(x) = 2^x$, and so $\varphi_g(x+1) = 2 * \varphi_g(x) = 2^{x+1}$ for all x , iff $\forall x \varphi_f(x) \downarrow$; otherwise $\varphi_g(x) \uparrow$ for some x .

Summarizing, f is in Total iff g is in DOUBLES and so

TOTAL \leq_m **DOUBLES** as we were to show.

Assignment # 8.4 Sample Key

4. Show **DOUBLES** reduces to **Total**. (3 plus 4 show they are equally hard)

Let f be an arbitrary natural number. f is in **DOUBLES** iff $\forall x \varphi_f(x) \downarrow$, $\varphi_f(x+1) \downarrow$ and $\varphi_f(x+1) = 2 * \varphi_f(x)$.

Define g by $\varphi_g(x) = \mu y [\varphi_f(x+1) = 2 * \varphi_f(x)]$, for all x .

Clearly, $\varphi_g(x) \downarrow$, for all x , iff $\forall x \varphi_f(x) \downarrow$, $\varphi_f(x+1) \downarrow$ and $\varphi_f(x+1) = 2 * \varphi_f(x)$; otherwise $\varphi_g(x) \uparrow$ for some x .

Summarizing, f is in **DOUBLES** iff g is in **Total** and so

DOUBLES \leq_m **TOTAL** as we were to show.

Assignment # 8.5 Sample Key

5. Use Rice's Theorem to show that **REPEATS** is undecidable

First, REPEATS is non-trivial as $C0(x) = 0$ is in REPEATS and $S(x) = x+1$ is not.

Second, REPEATS is an I/O property.

To see this, let f and g are two arbitrary indices such that

$$\forall x [\varphi_f(x) = \varphi_g(x)]$$

$f \in \text{REPEATS}$ iff $\exists y, z, y \neq z$, such that $\varphi_f(y) \downarrow, \varphi_f(z) \downarrow$ and $\varphi_f(y) = \varphi_f(z)$ iff, since $\forall x [\varphi_f(x) = \varphi_g(x)]$, $\exists y, z, y \neq z$, (same y, z as above) such that $\varphi_g(y) \downarrow, \varphi_g(z) \downarrow$ and $\varphi_g(y) = \varphi_g(z)$ iff $g \in \text{REPEATS}$.

Thus, **$f \in \text{REPEATS}$ iff $g \in \text{REPEATS}$** .

Assignment # 8.6 Sample Key

6. Use Rice's Theorem to show that **DOUBLES** is undecidable

First, DOUBLES is non-trivial as $C0(x) = 0$ ($2*0 = 0$) is in DOUBLES and $S(x) = x+1$ is not.

Second, DOUBLES is an I/O property.

To see this, let f and g are two arbitrary indices such that

$\forall x [\varphi_f(x) = \varphi_g(x)]$.

$f \in \text{DOUBLES}$ iff for all x , $\varphi_f(x) \downarrow$, $\varphi_f(x+1) \downarrow$ and $\varphi_f(x+1) = 2 * \varphi_f(x)$ iff, since $\forall x [\varphi_f(x) = \varphi_g(x)]$, for all x , $\varphi_g(x) \downarrow$, $\varphi_g(x+1) \downarrow$ and $\varphi_g(x+1) = 2 * \varphi_g(x)$ iff $g \in \text{DOUBLES}$.

Thus, **$f \in \text{DOUBLES}$ iff $g \in \text{DOUBLES}$.**