


# CAP 4453

## Robot Vision

Dr. Gonzalo Vaca-Castaño  
[gonzalo.vacacastano@ucf.edu](mailto:gonzalo.vacacastano@ucf.edu)






# Credits

- Slides comes directly from:
  - Ioannis (Yannis) Gkioulekas (CMU)
  - Noah Snavely (Cornell)
  - Marco Zuliani


# Short Review from last class





# Last 2 classes

- Feature points
  - Correspondent points on two images



# Robot Vision

## 9. Image warping I



# How do you create a panorama?

Panorama: an image of (near) 360° field of view.



# How do you create a panorama?

Panorama: an image of (near) 360° field of view.



1. Use a very wide-angle lens.



# Wide-angle lenses

Fish-eye lens: can produce (near) hemispherical field of view.



What are the pros and cons of this?



# How do you create a panorama?

Panorama: an image of (near) 360° field of view.



1. Use a very wide-angle lens.
  - Pros: Everything is done optically, single capture.
  - Cons: Lens is super expensive and bulky, lots of distortion (can be dealt-with in post).

Any alternative to this?

# How do you create a panorama?

Panorama: an image of (near) 360° field of view.



1. Use a very wide-angle lens.
  - Pros: Everything is done optically, single capture.
  - Cons: Lens is super expensive and bulky, lots of distortion (can be dealt-with in post).
2. Capture multiple images and combine them.



# Panoramas from image stitching

1. Capture multiple images from different viewpoints.



2. Stitch them together into a virtual wide-angle image.





# How do we stitch images from different viewpoints



Will standard stitching work?

1. Translate one image relative to another.
2. (Optionally) find an optimal seam.

# How do we stitch images from different viewpoints



Will standard stitching work?

1. Translate one image relative to another.
2. (Optionally) find an optimal seam.

left on top



right on top



Translation-only stitching is not enough to mosaic these images.

# How do we stitch images from different viewpoints



What else can we try?



# How do we stitch images from different viewpoints



Use image homographies.



# Outline

- Linear algebra
  - Matrix addition, Matrix multiplication
  - Inverse, Pseudo Inverse
  - Least squares, SVD
- Image transformations.
- 2D transformations.
- Projective geometry 101.
- Transformations in projective geometry.
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.

# Matrix

- Array  $\mathbf{A} \in \mathbb{R}^{m \times n}$  of numbers with shape m by n,
  - m rows and n columns

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- A row vector is a matrix with single row
- A column vector is a matrix with single column



# Matrix operations

- Addition

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a + e & b + f \\ c + g & d + h \end{bmatrix}$$


- Both matrices should have same shape, except with a scalar

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + 2 = \begin{bmatrix} a + 2 & b + 2 \\ c + 2 & d + 2 \end{bmatrix}$$

- Same with subtraction

# Matrix operation

- Matrix Multiplication
  - Compatibility?
  - $m \times n$  and  $n \times p$
  - Results in  $m \times p$  matrix



# Matrix operation

- Transpose

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$$\mathbf{A}^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}$$



# Special matrices

- Diagonal matrix
  - Used for row scaling

$$A = \begin{bmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{bmatrix}$$

- Identity matrix
  - Special diagonal matrix
  - 1 along diagonals

$$\mathbf{I} \cdot \mathbf{A} = \mathbf{A}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Matrix operation

- Inverse

- Given a matrix  $A$ , its inverse  $A^{-1}$  is a matrix such that

$$\mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$$

- Inverse does not always exist

- Singular vs non-singular

- Properties

- $(A^{-1})^{-1} = A$
  - $(AB)^{-1} = B^{-1}A^{-1}$

# Pseudoinverse

$$Ax = b \quad \leftarrow \text{A is not squared}$$

$$A^T Ax = A^t b \quad \leftarrow A^T A \text{ is squared}$$

$$(A^T A)^{-1}(A^T A)x = (A^T A)^{-1}A^t b$$

$$x = \underbrace{(A^T A)^{-1}A^t}_{\text{Pseudoinverse}} b$$



# Outline

- Linear algebra
- **Image transformations.**
- 2D transformations.
- Projective geometry 101.
- Transformations in projective geometry.
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.


# What is an image?

$$f(\mathbf{x})$$



grayscale image

What is the range of the image function  $f$ ?




domain  $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$


A (grayscale) image is a 2D function.

# Image Warping

- image filtering: change *range* of image




- image warping: change *domain* of image






# Image Warping

- image filtering: change *range* of image



- image warping: change *domain* of image



# What types of image transformations can we do




Filtering



changes pixel *values*



Warping



changes pixel *locations*

# What types of image transformations can we do

$F$



Filtering



$$G(\mathbf{x}) = h\{F(\mathbf{x})\}$$

$G$



changes *range* of image function

$F$




Warping



$$G(\mathbf{x}) = F(h\{\mathbf{x}\})$$

$G$



changes *domain* of image function



# The persistence of memory by Salvador Dali



Original



Filtering operation  
(Blurred)



Warping operation  
(Swirled)

What is the geometric relationship between these two images?



# What is the geometric relationship between these two images?



**Very important for creating mosaics!**

First, we need to know what this transformation is.

Second, we need to figure out how to compute it using feature matches.

# Warping example: feature matching






# Warping example: feature matching



# Warping example: feature matching




- object recognition
- 3D reconstruction
- augmented reality
- image stitching

How do you compute the transformation?

# Warping example: feature matching

Given a set of matched feature points:




and a transformation:

$$x' = f(x; p)$$

transformation function  $\nearrow$  parameters  $\nwarrow$

find the best estimate of the parameters

$p$



What kind of transformation functions  $f$  are there?

# Outline

- Linear algebra
- Image transformations
- **2D transformations.**
- Projective geometry 101.
- Transformations in projective geometry.
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.



# 2D transformations



translation



rotation



aspect



affine



perspective




cylindrical

# Parametric (global) warping



$\mathbf{p} = (x, y)$



$\mathbf{p}' = (x', y')$

- Transformation  $M$  is a coordinate-changing machine:

$$\mathbf{p}' = M(\mathbf{p})$$


- What does it mean that  $M$  is global?
  - Is the same for any point  $\mathbf{p}$
  - can be described by just a few numbers (parameters)
- Let's consider *linear* forms (can be represented by a 2x2 matrix):

$$\mathbf{p}' = M\mathbf{p} \qquad \begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{M} \begin{bmatrix} x \\ y \end{bmatrix}$$

# 2D planar transformations



# 2D planar transformations



$y$

Scale

How would you implement scaling?

- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

$x$



# 2D planar transformations

$y$



$$x' = ax$$

$$y' = by$$

What's the effect of using  
different scale factors?

- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

$x$

# 2D planar transformations

$y$



$$x' = ax$$

$$y' = by$$

matrix representation of scaling:


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \underbrace{\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}}_{\text{scaling matrix } S} \begin{bmatrix} x \\ y \end{bmatrix}$$

scaling matrix  $S$


- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

$x$

# 2D planar transformations




# 2D planar transformations







# 2D planar transformations



# 2D planar transformations



# 2D planar transformations



Polar coordinates...

$$x = r \cos(\phi)$$

$$y = r \sin(\phi)$$

$$x' = r \cos(\phi + \theta)$$

$$y' = r \sin(\phi + \theta)$$

Trigonometric Identity...

$$x' = r \cos(\phi) \cos(\theta) - r \sin(\phi) \sin(\theta)$$


$$y' = r \sin(\phi) \cos(\theta) + r \cos(\phi) \sin(\theta)$$

Substitute...

$$x' = x \cos(\theta) - y \sin(\theta)$$


$$y' = x \sin(\theta) + y \cos(\theta)$$

# 2D planar transformations



# Common linear transformations

- Rotation by angle  $\theta$  (about the origin)



$$\mathbf{R} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

What is the inverse?

For rotations:

$$\mathbf{R}^{-1} = \mathbf{R}^T$$



# 2D planar and linear transformations

$$\mathbf{x}' = f(\mathbf{x}; p)$$



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{M} \begin{bmatrix} x \\ y \end{bmatrix}$$

parameters  $p$

point  $\mathbf{x}$

# 2D planar and linear transformations

Scale

$$\mathbf{M} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}$$

Flip across y

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

Rotate

$$\mathbf{M} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

Flip across origin

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$


Shear

$$\mathbf{M} = \begin{bmatrix} 1 & s_x \\ s_y & 1 \end{bmatrix}$$

Identity

$$\mathbf{M} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

# 2D planar transformations



# All 2D Linear Transformations

- Linear transformations are combinations of ...

- Scale,
- Rotation,
- Shear, and
- Mirror


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Properties of linear transformations:

- Origin maps to origin
- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} \begin{bmatrix} i & j \\ k & l \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


# What is the geometric relationship between these two images?



**Answer: Similarity transformation** (translation, rotation, uniform scale)




# 2D translation




How would you implement translation?

# 2D translation



# 2D translation



# Outline

- Linear algebra
- Image transformations.
- 2D transformations.
- **Projective geometry**
- Transformations in projective geometry.
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.

# Homogeneous coordinates

heterogeneous  
coordinates

homogeneous  
coordinates

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

← add a 1 here

- Represent 2D point with a 3D vector



# Homogeneous coordinates


heterogeneous  
coordinates

homogeneous  
coordinates

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \stackrel{\text{def}}{=} \begin{bmatrix} ax \\ ay \\ a \end{bmatrix}$$

- Represent 2D point with a 3D vector
- 3D vectors are only defined up to scale

# 2D translation




$$x' = x + t_x$$

$$y' = y + t_y$$

What about matrix representation  
using homogeneous coordinates?

# 2D translation

$y$



$$x' = x + t_x$$

$$y' = y + t_y$$

What about matrix representation  
using heterogeneous coordinates?


$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \mathbf{M} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

$x$

# 2D translation using homogeneous coordinates



$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$



# Homogeneous coordinates

Conversion:

- heterogeneous  $\rightarrow$  homogeneous

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

- homogeneous  $\rightarrow$  heterogeneous

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

- scale invariance

$$\begin{bmatrix} x & y & w \end{bmatrix}^T = \lambda \begin{bmatrix} x & y & w \end{bmatrix}^T$$

Special points:

- point at infinity (x,y)

$$\begin{bmatrix} x & y & 0 \end{bmatrix}$$

- undefined

$$\begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$




# Homogeneous coordinates

Trick: add one more coordinate:

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image  
coordinates



Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

# Outline

- Linear algebra
- Image transformations.
- 2D transformations.
- Projective geometry
- **Transformations in projective geometry.**
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.

# 2D transformations in heterogeneous coordinates



Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

# 2D transformations in heterogeneous coordinates



Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

# 2D transformations in heterogeneous coordinates



Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & ? & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & \beta_x & 0 \\ \beta_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

# 2D transformations in heterogeneous coordinates



Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & \beta_x & 0 \\ \beta_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing



# Matrix composition

Transformations can be combined by matrix multiplication:

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \left( \begin{bmatrix} 1 & 0 & tx \\ 0 & 1 & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} sx & 0 & 0 \\ 0 & sy & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

$\mathbf{p}' = \quad ? \quad ? \quad ? \quad \mathbf{p}$

# Matrix composition

Transformations can be combined by matrix multiplication:

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \left( \begin{bmatrix} 1 & 0 & tx \\ 0 & 1 & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} sx & 0 & 0 \\ 0 & sy & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$


$\mathbf{p}'$  = translation( $t_x, t_y$ )      rotation( $\theta$ )      scale( $s, s$ )       $\mathbf{p}$

Does the multiplication order matter?

# Outline

- Linear algebra
- Image transformations.
- 2D transformations.
- Projective geometry
- Transformations in projective geometry.
- **Classification of 2D transformations.**
- Determining unknown 2D transformations.
- Determining unknown image warps.

# Classification of 2D transformations



# Classification of 2D transformations


Name	Matrix	# D.O.F.
translation	$\begin{bmatrix} \mathbf{I} &   & \mathbf{t} \end{bmatrix}$	?
rigid (Euclidean)	$\begin{bmatrix} \mathbf{R} &   & \mathbf{t} \end{bmatrix}$	?
similarity	$\begin{bmatrix} s\mathbf{R} &   & \mathbf{t} \end{bmatrix}$	?
affine	$\begin{bmatrix} \mathbf{A} \end{bmatrix}$	?
projective	$\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}$	?

# Classification of 2D transformations

Translation:

$$\begin{bmatrix} 1 & 0 & t_1 \\ 0 & 1 & t_2 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?






# Classification of 2D transformations

Euclidean (rigid):  
rotation + translation

$$\begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?




# Classification of 2D transformations

Euclidean (rigid):  
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?




# Classification of 2D transformations

which other matrix values will  
change if this increases?

Euclidean (rigid):  
rotation + translation

↓

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




# Classification of 2D transformations

what will happen to the  
image if this increases?

Euclidean (rigid):  
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




# Classification of 2D transformations

what will happen to the  
image if this increases?



Euclidean (rigid):  
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




# Classification of 2D transformations

Similarity:  
uniform scaling + rotation  
+ translation

$$\begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?





# Classification of 2D transformations


multiply these four by scale  $s$



Similarity:  
uniform scaling + rotation  
+ translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?




# Classification of 2D transformations

what will happen to the  
image if this increases?

Similarity:  
uniform scaling + rotation  
+ translation

$$\downarrow \begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




# Classification of 2D transformations

Affine transform:  
uniform scaling + shearing  
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

any transformation  
represented by a 3x3 matrix  
with last row  $[0 \ 0 \ 1]$  we call  
an *affine* transformation

Are there any values that are related?




# Classification of 2D transformations

Affine transform:  
uniform scaling + shearing  
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?

$$\begin{matrix} \text{similarity} \\ \begin{bmatrix} sr_1 & sr_2 \\ sr_3 & sr_4 \end{bmatrix} \end{matrix} \begin{matrix} \text{shear} \\ \begin{bmatrix} 1 & h_1 \\ h_2 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} sr_1 + h_2 sr_2 & sr_2 + h_1 sr_1 \\ sr_3 + h_2 sr_4 & sr_4 + h_1 sr_3 \end{bmatrix}$$




# Classification of 2D transformations

Affine transform:  
uniform scaling + shearing  
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?

$$\begin{matrix} \text{similarity} \\ \begin{bmatrix} sr_1 & sr_2 \\ sr_3 & sr_4 \end{bmatrix} \end{matrix} \begin{matrix} \text{shear} \\ \begin{bmatrix} 1 & h_1 \\ h_2 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} sr_1 + h_2 sr_2 & sr_2 + h_1 sr_1 \\ sr_3 + h_2 sr_4 & sr_4 + h_1 sr_3 \end{bmatrix}$$



# Affine transformations


Affine transformations are combinations of

- arbitrary (4-DOF) linear transformations; and
- translations

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of affine transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines map to parallel lines
- ratios are preserved
- compositions of affine transforms are also affine transforms



Does the last coordinate  $w$  ever change?



# Affine transformations


Affine transformations are combinations of

- arbitrary (4-DOF) linear transformations; and
- translations

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of affine transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines map to parallel lines
- ratios are preserved
- compositions of affine transforms are also affine transforms



Nope! But what does that mean?

# How to interpret affine transformations here?

image point in  
pixel coordinates

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$





image point in  
heterogeneous  
coordinates

$$\mathbf{X} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$



$\mathbf{X}$  is a projection of a point  
 $\mathbf{P}$  on the image plane

# Where do we go from here?

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$

affine transformation





what happens when we  
mess with this row?

# Projective Transformations aka Homographies aka Planar Perspective Maps

$$\mathbf{H} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix}$$

Called a *homography*  
(or *planar perspective map*)



# Homographies

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} \frac{ax+by+c}{gx+hy+1} \\ \frac{dx+ey+f}{gx+hy+1} \\ 1 \end{bmatrix}$$

# Alternate formulation for homographies

$$\begin{bmatrix} x'_i \\ y'_i \\ 1 \end{bmatrix} \cong \begin{bmatrix} h_{00} & h_{01} & h_{02} \\ h_{10} & h_{11} & h_{12} \\ h_{20} & h_{21} & h_{22} \end{bmatrix} \begin{bmatrix} x_i \\ y_i \\ 1 \end{bmatrix}$$

where the length of the vector  $[h_{00} \ h_{01} \ \dots \ h_{22}]$  is 1

# Projective transformations (aka homographies)

Projective transformations are combinations of


- affine transformations; and
- projective wraps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

How many degrees of freedom?

Properties of projective transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines do not necessarily map to parallel lines
- ratios are not necessarily preserved
- compositions of projective transforms are also projective transforms





# Projective transformations (aka homographies)

Projective transformations are combinations of


- affine transformations; and
- projective wraps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of projective transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines do not necessarily map to parallel lines
- ratios are not necessarily preserved
- compositions of projective transforms are also projective transforms

8 DOF: vectors (and therefore matrices) are defined up to scale)



# How to interpret projective transformations here?



image point in  
pixel coordinates

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$







image point in  
heterogeneous  
coordinates

$$\mathbf{X} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$




# Points at infinity








Is this an affine transformation?



# 2D image transformations



Name	Matrix	# D.O.F.	Preserves:	Icon
translation	$\begin{bmatrix} \mathbf{I} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	2	orientation + ...	
rigid (Euclidean)	$\begin{bmatrix} \mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	3	lengths + ...	
similarity	$\begin{bmatrix} s\mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	4	angles + ...	
affine	$\begin{bmatrix} \mathbf{A} \end{bmatrix}_{2 \times 3}$	6	parallelism + ...	
projective	$\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}_{3 \times 3}$	8	straight lines	

These transformations are a nested set of groups

- Closed under composition and inverse is a member



When can we use homographies?



# We can use homographies when...

1. ... the scene is planar; or



2. ... the scene is very far or has small (relative) depth variation  
→ scene is approximately planar





# We can use homographies when...

3. ... the scene is captured under camera rotation only (no translation or pose change)




More on why this is the case in a later lecture.




# Computing with homographies

# Classification of 2D transformations



Which kind transformation is needed to warp projective plane 1 into projective plane 2?

- A projective transformation (a.k.a. a homography).



# Applying a homography

1. Convert to homogeneous coordinates:

$$p = \begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow P = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

What is the size of the homography matrix? ↘

2. Multiply by the homography matrix:

$$P' = H \cdot P$$

3. Convert back to heterogeneous coordinates:

$$P' = \begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} \Rightarrow p' = \begin{bmatrix} x'/w' \\ y'/w' \end{bmatrix}$$

# Applying a homography

1. Convert to homogeneous coordinates:  
$$p = \begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow P = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$
2. Multiply by the homography matrix:  
$$P' = H \cdot P$$

What is the size of the homography matrix?  $\searrow$  Answer: 3 x 3

How many degrees of freedom does the homography matrix have?  $\nearrow$
3. Convert back to heterogeneous coordinates:  
$$P' = \begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} \Rightarrow p' = \begin{bmatrix} x'/w' \\ y'/w' \end{bmatrix}$$

# Applying a homography

1. Convert to homogeneous coordinates:  
$$p = \begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow P = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$
2. Multiply by the homography matrix:  
$$P' = H \cdot P$$

What is the size of the homography matrix?  $\searrow$  Answer: 3 x 3

How many degrees of freedom does the homography matrix have?  $\nearrow$  Answer: 8
3. Convert back to heterogeneous coordinates:  
$$P' = \begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} \Rightarrow p' = \begin{bmatrix} x'/w' \\ y'/w' \end{bmatrix}$$



# Applying a homography

What is the size of the homography matrix?

Answer: 3 x 3



$$P' = H \cdot P$$



How many degrees of freedom does the homography matrix have?

Answer: 8

How do we compute the homography matrix?



# Homography

Under homography, we can write the transformation of points in 3D from camera 1 to camera 2 as:


$$\mathbf{X}_2 = H\mathbf{X}_1 \quad \mathbf{X}_1, \mathbf{X}_2 \in \mathbb{R}^3 \quad \leftarrow \text{Homogeneous coordinates}$$

In the image planes, using homogeneous coordinates, we have

$$\lambda_1 \mathbf{x}_1 = \mathbf{X}_1, \quad \lambda_2 \mathbf{x}_2 = \mathbf{X}_2, \quad \text{therefore} \quad \lambda_2 \mathbf{x}_2 = H \lambda_1 \mathbf{x}_1$$

$\lambda_1 \mathbf{x}_1$   $\lambda_2 \mathbf{x}_2$   
Heterogeneous coordinates

**This means that  $\mathbf{x}_2$  is equal to  $H\mathbf{x}_1$  up to a scale (due to universal scale ambiguity)**



# Outline

- Linear algebra
- Image transformations
- 2D transformations.
- Projective geometry 101.
- Transformations in projective geometry.
- Classification of 2D transformations.
- **Determining unknown 2D transformations.**
- Determining unknown image warps.



# References

Basic reading:

- Szeliski textbook, Section 3.6.

Additional reading:

- Hartley and Zisserman, "Multiple View Geometry in Computer Vision," Cambridge University Press 2004.  
a comprehensive treatment of all aspects of projective geometry relating to computer vision, and also a very useful reference for the second part of the class.
- Richter-Gebert, "Perspectives on projective geometry," Springer 2011.  
a beautiful, thorough, and very accessible mathematics textbook on projective geometry (available online for free from CMU's library).



# Questions?